



Subscribe | Login | Register | Claim Subscription

Home Articles & Issues Editor's Choice For Authors Journal Info Subscribe EPA More Periodicals Contact

All Content

Search Advanced Search

Previous Article
2011 Volume 26, Supplement 1, Page 1463

Next Article >

Po3-294 - Severe generalize cortical atrophy inconsistent with her age, and neurocognitive impairment in an untreated schizophrenia patient: A case report

E. Oral, M. Güleç, N. Aydin, E. Ozan, I. Kırpınar

Psychiatry, Ataturk University, Faculty of Medicine, Erzurum, Turkey



DOI: https://doi.org/10.1016/S0924-9338(11)73168-6

Abstract

Background

In schizophrenia, frontal structural and functional deficits are replicated findings.

Case

AK was 40 years old. She had not been appealed to psychiatry department or took any medicine. She had had severe positive symptoms such as, grandiosity, delusions, auditory and visual hallucinations, hallucinatory behaviors, mystic contains of thoughts since her childhood. MRI had been showed that the severe cortical atrophy inconsistent with her age. In the assessment of neurocognitive tests we found that the signs of frontal -prefrontal and temporo-limbic impairment. On Rey Auditory learning test she could remember totally 34 words as a sum of the 1st to 5.th Readings. At 6.th reading she remembered 5 words and at 7th 6 words. At recognition part of the test she had 8 true positive and 3 false positive results. On Auditory trigram test, her score was 25 and on controlled word association test she could tell 20 total words. In digit span test her scores were 4 at forward section and 4 at backwards section. She could not perform trial making test part A or part B. On Stroop test she completed section 5 (main card reading time) at 26 seconds. These results suggested us that she had dysfunction in the cognitive domains such as; working memory, focused attention, short time verbal memory and frontal executive functions.

Conclusion

This sample has been attracted attention to neurodegenerative process notwithstanding without antipsychotic treatments effects.

© 2011 Elsevier Masson SAS. Published by Elsevier Inc. All rights reserved.

Access this article on ScienceDirect

Article Tools

PDF (12 KB)

Email Article

Add to My Reading List

Export Citation

Create Citation Alert

Cited by in Scopus (0)

Request Permissions

Order Reprints
(100 minimum order)

Related Articles

Impaired self-agency inferences in schizophrenia: The role of cognitive capacity and causal reasoning style European Psychiatry, Vol. 47

Daily life impairments associated with self-reported childhood/adolescent attention-deficit/hyperactivity disorder and experiences of diagnosis and treatment: Results from the European Lifetime Impairment Survey
European Psychiatry, Vol. 29, Issue 5

Six-year follow-up study of combined type ADHD from childhood to young adulthood: Predictors of functional impairment and comorbid symptoms European Psychiatry, Vol. 35

Prediction of motivational impairment: 12month follow-up of the randomizedcontrolled trial on extended early intervention for first-episode psychosis European Psychiatry, Vol. 41

Impaired action self-monitoring and cognitive confidence among ultra-high risk for psychosis and first-episode psychosis patients European Psychiatry, Vol. 47

View All

< Previous Article

2011 Volume 26, Supplement 1, Page 1463

Next Article >

Copyright © 2018 Elsevier Inc. All rights reserved. | Privacy Policy | Terms & Conditions | Use of Cookies | About Us | Help & Contact | Accessibility The content on this site is intended for health professionals.

Advertisements on this site do not constitute a guarantee or endorsement by the journal, Association, or publisher of the quality or value of such product or of the claims made for it by its manufacturer.

04.02.2018	P03-294 -	- Severe generalize cortical atrophy inconsistent with her age, and neurocognitive impairment in an untreated schizophrenia pati